

Nonlinear electrodynamics in Bianchi spacetimes

Ricardo García-Salcedo*

*Departamento de Física, Cinvestav-IPN, Apartado Postal 14-740, C.P. 07000, México, D.F., MEXICO and
Facultad de Ciencias, U. A. E. Mex., Instituto Literario 100, 50000, Toluca, México.*

Nora Bretón†

*Departamento de Física, Cinvestav-IPN, Apartado Postal 14-740, C.P. 07000, México, D.F., MEXICO
(Dated: February 1, 2008)*

We study the effects produced by nonlinear electrodynamics in spacetimes conformal to Bianchi metrics. In the presence of Born-Infeld type fields these models accelerate, expand and isotropize. This effect is compared with the corresponding to a linear electromagnetic field; it turns out that for the same geometry, Maxwell field does not favour inflation as much as Born-Infeld field. The behavior of the nonlinear radiation is analyzed in terms of the equations of state. The energy conditions are analyzed as well, showing that the Born-Infeld field violates the strong energy condition.

PACS numbers: 98.80.Hw, 11.10.Lm, 12.20.-m

I. INTRODUCTION

Born and Infeld proposed their theory for nonlinear electrodynamics (NLE) in 1934 [1] with the aim of removing the point charge singularity in classical electrodynamics. However, soon it became clear that the Born-Infeld (BI) theory works as well as a phenomenological theory of quantum electrodynamics in the first approximation. The effective action for electrodynamics due to one-loop quantum corrections was calculated by Heisenberg and Euler [2], for the low frequency limit; the corresponding effective Lagrangian have quadratic terms in the electromagnetic field invariants like the BI Lagrangian. Moreover, actions of the BI type have recently been the subject of wide interest in relation to the result that the effective action for the open string ending on D-branes can be written in a BI form. In this respect, BI cosmological solutions could assist in the understanding of brane dynamics regarding the universe evolution.

These facts make NLE particularly interesting in general relativity, specially in cosmological theories, as a simple classical model that incorporates vacuum polarization processes -a possible influence on the mechanism of the evolution of the early universe; these processes are expressed in terms of a kind of electric permittivity and magnetic permeability of vacuum. From this point of view, the exact solutions of Einstein-Born-Infeld (EBI) equations are worth to study since they may indicate the physical relevance of nonlinear effects in strong gravitational and strong electromagnetic fields at early epochs of the universe. Moreover, in [3] it was found that BI cosmological models can be compatible with the recently measured accelerated expansion of the universe.

One of the topics of interest is to elucidate to what extent NLE favours or not an inflationary scenario. The

cosmological inflationary scenario, as was proposed by Guth [4], has been much favoured due to its troubleshooting capabilities. Furthermore, it has recently gained the support of observational evidence [5]. The conventional models use some special scalar fields with associated potential to give rise inflation. Another proposal to produce inflation is to use theories with unconventional dependence of the Lagrangian on field derivatives [6]. In this paper non-linear electrodynamics will be considered as the dynamical field which drives inflation.

On the other hand, there is no reason to assume that at the onset of inflation the universe was as isotropic as it seems to be today. It may be precisely an epoch of accelerated expansion that isotropized the universe. The most simple anisotropic models are the Bianchi ones. Inflation in Bianchi models have been addressed in the Einstein-Cartan theory and with a cosmological constant [7]. Inflation with a Brans-Dicke field in Bianchi V has been studied in [8]; assisted inflation was investigated in Bianchi I in [9] and in Bianchi VI_0 in [10].

In [11] an exact solution of the coupled Einstein-Born-Infeld equations for anisotropic cosmological scenario was presented. Later, other cosmological models have been proposed with nonlinear electromagnetic sources: in [12] it was obtained a homogeneous and isotropic nonsingular Friedman-Robertson-Walker (FRW) solution considering a nonlinear generalization of Maxwell electrodynamics as the source; in [13] it was investigated a Yang-Mills cosmology with non-Abelian BI action.

Here we present a study of the effects produced by a BI type field in an anisotropic cosmology, based on the analysis of the behavior of the mean Hubble parameter and the deceleration as well as the energy conditions. It turns out that inflation is produced, and in fact, for certain values of the BI parameter, the inflation is enough to provide the desired e-folding number that corresponds to the expansion that solves the cosmological problem of inflation. The model isotropizes at large times, in such a way that the entry and exit of inflation arise naturally as a consequence of dynamics. We also present the study of

*Electronic address: rsalcedo@fis.cinvestav.mx

†Electronic address: nora@fis.cinvestav.mx

the equations of state of the nonlinear radiation.

The plan of the paper is as follows: In Sec. II the nonlinear electrodynamics Lagrangian is introduced. In Sec. III the studied metric is presented; in Sec. IV it is given the corresponding group structure of the Killing vectors and it is shown that the studied space is conformal to the Bianchi spaces I, II, III, VII, VIII, IX, for particular values of the parameters. In section V the kinematical parameter are analyzed as functions of the synchronous time; comments are given on the e-folding number; in Sec. VI the effect produced by a linear electromagnetic field is analyzed. In Sec VII the pressure and energy density of the nonlinear electromagnetic matter is studied as well as the energy conditions. In Sec. VIII final remarks are given.

II. NONLINEAR ELECTRODYNAMICS THEORY

The nonlinear electromagnetic theory proposed by Born and Infeld is self-consistent and satisfies all natural requirements. Its Lagrangian depends in nonlinear form on the invariants of the field,

$$L_{BI} = -\frac{1}{2}P^{\mu\nu}F_{\mu\nu} + \mathcal{H}(P, Q), \quad (1)$$

where $\mathcal{H}(P, Q)$ is the so called structural function; this function can vary from one version of nonlinear electrodynamics to other. P and Q are the invariants associated with the antisymmetric tensor $P_{\mu\nu}$ that is the generalization of the electromagnetic tensor $F_{\mu\nu}$. The structural function is constrained to satisfy some physical requirements: (i) the correspondence to the linear theory [$\mathcal{H}(P, Q) = P + O(P^2, Q^2)$]; (ii) the parity conservation [$\mathcal{H}(P, Q) = \mathcal{H}(P, -Q)$]; (iii) the positive definiteness of the energy density ($\mathcal{H}_{,P} > 0$) and the requirement of the timelike nature of the energy flux vector ($P\mathcal{H}_{,P} + Q\mathcal{H}_{,Q} - \mathcal{H} \leq 0$). Condition (iii) amounts to the fulfilment of the dominant energy condition (DEC) stated as [14]: $T_{\mu\nu}u^\mu u^\nu > 0$ for every time direction u_μ , ($u^\mu u_\mu < 0$). The energy-momentum tensor is given by:

$$T_{\mu\nu} = -\mathcal{H}_{,P}P_{\mu\alpha}P^\alpha_\nu + g_{\mu\nu}[2P\mathcal{H}_{,P} + Q\mathcal{H}_{,Q} - \mathcal{H}]. \quad (2)$$

The structural function for the BI field is given by

$$\mathcal{H} = b^2 - \sqrt{b^4 - 2b^2P + Q^2}, \quad (3)$$

where b is the Born-Infeld constant that is the “absolute or maximum field”; this parameter (comparable to a coupling constant) distinguishes different theories. For the problem of the electron the value of the constant is found to be $b = e/r_0^2 = 9.18 \times 10^{15}$ e.s.u., while for the Heisenberg-Euler theory $b = 14\alpha^2/45m^4$, where α is the fine structure constant and m is the mass of the electron;

in string theory, $b = 2\pi\alpha'$ where α' is the string tension parameter. Here we shall consider that b is a parameter that can vary in order to determine if Lagrangians of the type (1) are consistent with an inflationary behavior with a b tuned for that purpose. The linear limit is obtained by taking $b \rightarrow \infty$, then $\mathcal{H} = P$ and $P_{\mu\nu} = F_{\mu\nu}$, coinciding with the Maxwell electromagnetism.

$P_{\mu\nu}$ and $F_{\mu\nu}$ are related through the material or constitutive equations:

$$F_{\mu\nu} = \mathcal{H}_{,P}P_{\mu\nu} + \mathcal{H}_{,Q}\check{P}_{\mu\nu}, \quad (4)$$

where $\check{P}_{\mu\nu} = -\frac{1}{2}\epsilon_{\mu\nu\alpha\beta}P^{\alpha\beta}$ is the dual of $P_{\mu\nu}$. An extensive treatment on nonlinear electrodynamics can be found in [15].

III. EBI SOLUTION IN AN ANISOTROPIC SPACETIME

We can think of a universe filled with radiation governed by non-linear dynamics. In [11] it was presented an exact solution to the coupled system of EBI equations for an anisotropic cosmological model without further investigations on kinematical aspects nor inflation. The spacetime is given by the line element

$$ds^2 = \frac{1}{\varphi^2} \left\{ \frac{dz^2}{h} - \frac{dt^2}{s} + hdy^2 + s(dx + Mdy)^2 \right\}, \quad (5)$$

where x, y are ignorable coordinates, $h = \alpha + \epsilon z^2$, ϵ, α and l are constants, $s = s(t)$, $\varphi = \varphi(t)$ and $M = 2lz$. The motivation to choose a metric of the form (5) is that it is a type D metric which for the stationary axisymmetric case (one spacelike Killing vector and one timelike Killing vector) solutions are known of coupled Einstein-Born-Infeld fields [16]. We expected, on the other side, that metric (5) be related to G_2 cosmologies, however the analysis of the Killing vectors shows (see next Section) that metric (5) possesses four Killing vectors and that it is conformal to Bianchi metrics. The spacetime (5) is coupled to an NLE energy-momentum tensor given by Eqs. (2) and (3), such that the two nonvanishing components of the electromagnetic field have been aligned with the two double repeated directions of the Weyl tensor and parametrized by $P_{12} = i\check{H}$, $P_{34} = D$, where $\check{H}$ is the magnetic field and D is the electric displacement; the invariants of the BI field are then given by $P = -\frac{1}{2}(D^2 - \check{H}^2)$ and $Q = -i\check{H}D$. We introduce the function $\nu(t)$ defined by $e^\nu = \sqrt{b^2 - \check{H}^2}/\sqrt{b^2 + D^2}$.

The solution to the coupled EBI equations is given by the metric function $s(t)$ in terms of $\varphi(t)$ (see [11] for details),

$$s(t) = \varphi^3 \dot{\varphi} \left\{ c_1 - 2b^2 \int \frac{(e^\nu - 1)}{\varphi^4 \dot{\varphi}^2} dt - \epsilon \int \frac{dt}{\varphi^2 \dot{\varphi}^2} \right\}, \quad (6)$$

where dot simbolizes derivative with respect to t , c_1 is an integration constant related to the solution in vacuum, the second term corresponds to the BI contribution and the last term is related to the spatial curvature, being $\epsilon = 1$ for a closed universe, $\epsilon = 0$ for spatially flat universe and $\epsilon = -1$ for an open universe. Moreover, $\varphi(t)$ and $\nu(t)$ must satisfy the equations $\ddot{\varphi} + l^2\varphi = 0$ and $\dot{\nu}\varphi - 2\dot{\varphi}(1 - e^{-2\nu}) = 0$.

The system of EBI equations admits two possible solutions: $e^\nu = \sqrt{1 \pm \varphi^4}$. The case $e^\nu = \sqrt{1 - \varphi^4}$ is the one that admits a cosmological interpretation, since in this case $s(t) > 0$. The case $e^\nu = \sqrt{1 + \varphi^4}$ corresponds to a metric function $s(t) < 0$ which changes the signature of the metric (5) to a stationary axisymmetric one. In this work we restrict to the range of t for which $s(t) > 0$; otherwise the signature of the metric changes and another interpretation must be considered.

The condition that e^ν be real imposes the restriction that $\varphi^4 < 1$; in terms of the fields it amounts to $\dot{H} < b$ and $D < b$, that is, the fields must not reach the maximum field strenght b . Also this condition determines the range of the coordinate t with $0 \leq \varphi(t) < 1$.

Moreover, the only nonvanishing Weyl scalar is given by

$$\psi_2 = \frac{\varphi^2}{12}[\ddot{s} - 8sl^2 - 2\epsilon + 6il\dot{s}]. \quad (7)$$

The invariants of this spacetime are $I = 3\psi_2^2$, $J = -\psi_2^3$, in such a manner that if ψ_2 diverges, so do the invariants. For this solution, ψ_2 is well behaved and does not present divergences. Up to here these results were derived in [11] and what follows is the contribution of the present paper.

IV. CHARACTERIZATION OF THE SPACETIME

In this section it is shown that the metric (5) is conformal to Bianchi spaces I, II, III, VII, VIII and IX.

The metric (5) possesses four Killing vectors:

$$\begin{aligned} \xi_1 &= [0, 1, 0, 0], & \xi_2 &= [0, 0, 1, 0], \\ \xi_3 &= [\sqrt{h}S(\sqrt{\epsilon\alpha}y), -\frac{2laC(\sqrt{\epsilon\alpha}y)}{\sqrt{h\epsilon\alpha}}, -\frac{\epsilon zC(\sqrt{\epsilon\alpha}y)}{\sqrt{h\epsilon\alpha}}, 0], \\ \xi_4 &= [\sqrt{h}C(\sqrt{\epsilon\alpha}y), -\frac{2laS(\sqrt{\epsilon\alpha}y)}{\sqrt{h\epsilon\alpha}}, -\frac{\epsilon zS(\sqrt{\epsilon\alpha}y)}{\sqrt{h\epsilon\alpha}}, 0] \end{aligned} \quad (8)$$

where S stands for $\sinh$ and C for $\cosh$. This set corresponds to a four-generator group with invariance subgroups of spatial rotations or boosts. From the above expressions is easy to show the commutation rules, $[\xi_1, \xi_3] = \sqrt{\epsilon\alpha}\xi_4$; $[\xi_1, \xi_4] = \sqrt{\epsilon\alpha}\xi_3$; $[\xi_3, \xi_4] = \sqrt{\epsilon\alpha}\xi_1$.

The metric (5) is conformal to several Bianchi spaces,

TABLE I: Values of α and ϵ for which the studied metric is conformal to the Bianchi types

σ^1	$h(y)$	α	ϵ	Bianchi type
$dx + 2l \cos y dz$	$\sin^2 y$	1	-1	G ₃ IX
$dx + 2ly dz$	1	1	0	G ₃ II
$dx + 2l \cosh y dz$	$\sinh^2 y$	-1	1	G ₃ III, G ₃ VIII

it is demonstrated by performing the transformation

$$\frac{dz^2}{h(z)} = dy^2, \quad \frac{dt^2}{s(t)} = dT^2, \quad y = z, \quad x = x, \quad (9)$$

with this transformation the metric, Eq. (5), acquires the form

$$\varphi^2 ds^2 = -dT^2 + s(T)(\sigma^1)^2 + dy^2 + h(y)dz^2, \quad (10)$$

where we named $dx + M(y)dz = \sigma^1$. Bianchi types IX, II, III and VIII can be identified according to Table I (cf. Eq. (11.4) in [17]).

Furthermore, in the case $M = 0$, ($l = 0$) the line element (5) becomes conformal to G₃I or G₃VII for $\epsilon = 0$ and G₃III for $\epsilon = 1$.

Also, with a coordinate transformation, the metric (5) can be identified as a hypersurface-homogeneous spacetime with a four dimensional group of isometries, G_4 , with three dimensional subgroups that can be of spatial rotations or boosts (S_3 or T_3),

$$\varphi^2 ds^2 = -\frac{4d\zeta d\bar{\zeta}}{(1 + \epsilon\zeta\bar{\zeta})^2} - \frac{dt^2}{s(t)} + s(t) \left(dx + 2il \frac{\zeta d\bar{\zeta} - \bar{\zeta} d\zeta}{(1 + \epsilon\zeta\bar{\zeta})} \right)^2, \quad (11)$$

The needed transformation to pass from Eq. (5) to Eq.(11) is (t remains the same):

$$z = -a \frac{(1 - \epsilon\zeta\bar{\zeta})}{(1 + \epsilon\zeta\bar{\zeta})}, \quad y = \frac{i}{2a\epsilon} \ln \frac{\bar{\zeta}}{\zeta}, \quad x = x + \frac{il}{\epsilon} \ln \frac{\bar{\zeta}}{\zeta}. \quad (12)$$

In the form (11) it is manifest the existence of a two-dimensional submanifold of constant curvature.

V. INFLATION PRODUCED BY THE BI FIELD

In this section we study the kinematical behavior in the asymptotic regimes of $\varphi \rightarrow 1$ and $\varphi \rightarrow 0$, that correspond to $t \rightarrow 0$ and $t \rightarrow \pi/2$, respectively; in a synchronous reference system with synchronous time given by $dT = \frac{dt}{\varphi\sqrt{s}}$, $\varphi \rightarrow 1$ and $\varphi \rightarrow 0$ correspond to $T \rightarrow 0$ and $T \rightarrow \infty$, respectively. In this analysis we shall restrict to the effect produced by the BI field, meaning that in the metric function $s(t)$, Eq. (6), we choose $c_1 = 0 = \epsilon$, then

$$s(t) = -2b^2\varphi^3\dot{\varphi} \int \frac{(e^\nu - 1)}{\varphi^4\dot{\varphi}^2} dt = b^2 S_{BI} \quad (13)$$

Moreover, we approximate $e^\nu - 1 = \sqrt{1 - \varphi^4} - 1 \approx -\varphi^4(1 + \varphi^4/4 + \varphi^8/8)/2$, approximation valid since $\varphi^4 < 1$. Then in the asymptotic regimes the limits for the metric function $s(t)$ and its derivative are

$$\begin{aligned} s(t \rightarrow 0) &\rightarrow 11b^2/8; & \dot{s}(t \rightarrow 0) &\rightarrow 0. \\ s(t \rightarrow \pi/2) &\rightarrow 0; & \dot{s}(t \rightarrow \pi/2) &\rightarrow 0. \end{aligned} \quad (14)$$

We now proceed to analyze the Hubble, deceleration and shear parameters. In the synchronous reference system where $g_{00} = 1$ and $g_{0i} = 0$, we introduce a spatial metric tensor h_{ab} . The volume element in three space is $\sqrt{h} = \varphi^{-3}\sqrt{s}$, ($h = \det h_{ab}$), while the volume expansion or mean Hubble parameter is given by

$$H = \frac{\varphi\sqrt{s}}{6} \left(\frac{\dot{s}}{s} - 6\frac{\dot{\varphi}}{\varphi} \right), \quad (15)$$

The Hubble parameter has been calculated using the timelike unit vector, $U^\alpha = \varphi\sqrt{s(t)}\delta_t^\alpha$, or in the null tetrad formalism, $U^1 = U^2 = 0$ and $U^3 = -U^4 = \frac{1}{\sqrt{2}}$. In Eq. (15) H is a function of t . The role of the constant c_1 in the metric function $s(t)$, Eq.(6), when substituted in (15) is to shift the time for which the expansion becomes null. The instant when $H = 0$ can be placed before, at or after $t = 0$ by changing the value of c_1 .

In Fig. 1 it is shown a parametric plot of the mean Hubble parameter $H(T)$ in terms of the synchronous time. $H(T)$ is plotted for several values of the Born-Infeld parameter b . It is clear that as greater is b , more expansion is produced. After some time $H(T)$ goes to a constant. To generate the graphics we have chosen $\varphi = A \cos(lt + \beta)$ as the solution of $\ddot{\varphi} + l^2\varphi = 0$ and we have put the phase to zero.

The expression of the Hubble parameter considering only the BI contribution clearly shows its b dependence,

$$H_{BI} = b \frac{\varphi\sqrt{S_{BI}}}{6} \left(\frac{\dot{S}_{BI}}{S_{BI}} - 6\frac{\dot{\varphi}}{\varphi} \right). \quad (16)$$

The analysis of H_{BI} in the asymptotic regimes shows that both at $t \rightarrow 0$ ($T \rightarrow 0$) and at $t \rightarrow \pi/2$ ($T \rightarrow \infty$) $H_{BI} \rightarrow 0$, so the expansion occurs during a limited period of time.

The behavior of the Hubble parameter is not enough to determine whether the model inflates or not, rather we must determine the sign of the deceleration parameter, and it must be negative if inflation is going to happen. The deceleration is $q = -\theta^{-2}(3\theta_\alpha U^\alpha + \theta^{-2})$, where θ is the expansion of a timelike vector; in terms of $s(t)$ and $\varphi(t)$ the deceleration is given by

$$q = 3\theta^{-2}(\dot{\varphi}\varphi\dot{s} + \varphi^2(\dot{s}^2/4s - \ddot{s}/2 - 3l^2s)) - 1. \quad (17)$$

Using the field equation $2b^2(e^{-\nu} - 1) = \varphi^2\ddot{s}/2 - 2\dot{\varphi}\varphi\dot{s} + 3s(\dot{\varphi}^2 + l^2\varphi^2)$ and approximating $e^{-\nu} \approx 1 + \varphi^4/2 + 3\varphi^8/8$,




FIG. 1: It is shown the mean Hubble parameter H as function of the synchronous time T for distinct values of the BI parameter b (shown in parenthesis). It can be observed that the expansion has a maximum and after some time decays to a constant value. In this plot $\varphi = .8 \cos t$, $\epsilon = 0$, $c_1 = 0$.

which is valid since $\varphi^4 < 1$ and considering only the BI effect we obtain

$$q = \frac{b^2}{\theta^2} \left\{ \frac{\varphi^2 \dot{S}_{BI}^2}{2S_{BI}} - 3\varphi^4 \left(1 + \frac{3}{4}\varphi^4 + \frac{5}{8}\varphi^8 \right) \right\}. \quad (18)$$

The sign of the deceleration parameter is determined by the term in curly brackets, it is the second term, which can make q to be negative, since S_{BI} , φ^2 and $\dot{S}_{BI}^2$ are positive. It is also illustrative that the asymptotic behavior of q as t tends to zero is $q(t \rightarrow 0) \rightarrow -\infty$; the infinite is due to the fact that as t tends to zero the expansion θ also tends to vanish; generically for the BI contribution we can say that q is negative as the time tends to zero. Regarding the asymptotic regime of $t \rightarrow \pi/2$ ($T \rightarrow \infty$), q tends to vanish, $q(t \rightarrow \pi/2) \rightarrow 0$, with this showing that the deceleration stops as the synchronous time goes to infinity. Fig. 2 shows the behavior of q , the plot corresponds to $q\theta^2/b^2$.

The behavior described above of the expansion and deceleration parameters is qualitatively the same for the cases $\epsilon = 1, 0, -1$ (different Bianchi spaces). For distinct values of ϵ the inflationary period changes its duration: for $\epsilon = -1$, q is negative but greater in absolute value during a shorter period. For $\epsilon = 1$, q is negative but smaller in absolute value than for $\epsilon = -1$, but q is negative during a longer period. The point where q diverges to negative infinity is the same for which $H = 0$, then in this respect the shifting in time also applies for q : by changing the value of c_1 in the function $s(t)$, the value of t for which q diverges is shifted.

It has been shown that there exists an accelerated expansion produced by the BI type field. How much does inflate a spacetime is specified by the e-folding number N . The cosmological problem of expansion is solved with




FIG. 2: The qualitative behavior of the deceleration parameter q is displayed in this figure: at the beginning there is acceleration followed by a deceleration period, $q > 0$, and finally, as $t \rightarrow \pi/2$, $q \rightarrow 0$. The plot corresponds to $q\theta^2/b^2$ and $s(t) = b^2 S_{BI}$, with $\varphi = \cos t$.




FIG. 3: It is shown a parametric plot of the shear σ and the synchronous time T , for several values of b (shown in parenthesis). The model isotropizes after some time and the maximum of the shear is higher for larger values of the parameter b . Here $\varphi = .8 \cos t$, $c_1 = 0$ and $\epsilon = 0$.

a minimum e-folding number of about 70, that amounts to an expansion of 10^{30} . The number of e-folds of growth in the scale factor that occurs from t_1 to t_2 is given by

$$N = \int_{t_1}^{t_2} H_{BI} dt = b \int_{t_1}^{t_2} \frac{\varphi \sqrt{S_{BI}}}{6} \left(\frac{\dot{S}_{BI}}{S_{BI}} - 6 \frac{\dot{\varphi}}{\varphi} \right) dt, \quad (19)$$

Since the integrand is positive the number $N \sim 70$ can be reached by adjusting the parameter b ; for $b \sim 10^3$ it is obtained the desired N .

In relation with the isotropy of the model, it has been shown within the framework of general relativity [18] that if the dominant energy conditions and positive pressure criterion are satisfied then the universe will isotropize




FIG. 4: The expansion H , deceleration q and the metric function $s(t)$ produced by the linear electromagnetic field are shown in this figure. In this case inflation occurs during a short period of time which ends when the deceleration becomes positive (at $t = 0.32$). Here $\varphi = \cos t$, $C = 1$, $c_1 = 0$ and $\epsilon = 0$.

starting from Bianchi I, V, VII_o, and VII_h models. The anisotropic character is given by the shear, σ , then the necessary condition for eventually obtain a FRW cosmology is the vanishing of the shear. For our model the shear is given by $\sigma^2 = \varphi^2 \dot{s}^2 / 6s$. Then a requirement for the model to isotropize is that the derivative of $s(t)$ goes to zero after some time. The tendency of the shear at the asymptotic regimes is such that it vanishes both as t goes to zero and as t approximates $\pi/2$, $\sigma(t \rightarrow 0) \rightarrow 0$ and $\sigma(t \rightarrow \pi/2) \rightarrow 0$. Then the shear grows at early times in the model and as $t \rightarrow \pi/2$ ($T \rightarrow \infty$ in the synchronous time) it gradually decays reaching zero as the synchronous time goes to infinity. In Fig. 3 the shear is shown in terms of the synchronous time T , for several values of the BI field; the peak of σ is greater as larger is the parameter b . This plot shows how these BI cosmologies isotropizes. Only the BI effect is considered in the graphics, meaning by this that the constants in $s(t)$, Eq (6), are $c_1 = 0 = \epsilon$.

VI. INFLATION PRODUCED BY A LINEAR ELECTROMAGNETIC FIELD

It is interesting to compare the effects produced by the BI field with those corresponding to the limit of linear electromagnetic field. This effect is obtained by taking $b \rightarrow \infty$. The process of taking this limit involves at the same time to take it in the function $e^\nu = \sqrt{b^2 - \check{H}^2} / \sqrt{b^2 + D^2}$, in such a manner that when $b \rightarrow \infty$ we have

$$\begin{aligned} 2b^2(e^{-\nu} - 1) &\rightarrow D^2 + \check{H}^2, \\ -2b^2(e^\nu - 1) &\rightarrow D^2 + \check{H}^2. \end{aligned} \quad (20)$$




FIG. 5: In this plot it is shown the two distinct eigenvalues of T^{ab} : $\rho = p_1 = T_{34}$ and $p = p_2 = p_3 = T_{12}$. ρ is the energy density and p_1, p_2, p_3 are the pressures in the three spatial directions that measures an observer at the point p^* , whose world line has as tangent the timelike vector E_4 . The observer measures in one direction a stiff fluid of positive pressure ($p = \rho$) and in the other two directions there is a kind of “vacuum” state with negative pressures ($p_2 = p_3 < 0$).

Using this limits in the expressions for the metric function $s(t)$ as well as in the expansion and deceleration we obtain the corresponding quantities when a linear electromagnetic field is acting on the metric (5). They are:

$$s(t) = \varphi^3 \dot{\varphi} \left\{ c_1 + C^2 \int \frac{dt}{\varphi^4 \dot{\varphi}^2} - \epsilon \int \frac{dt}{\varphi^2 \dot{\varphi}^2} \right\}, \quad (21)$$

$$q = \frac{2\dot{s}^2 \varphi^2 - 12sC^2}{(6s\dot{\varphi} - \dot{s}\varphi)^2}, \quad (22)$$

where $C^2 = D^2 + \ddot{H}^2$ is related with the energy density of the linear electromagnetic field; the expansion H is given as in Eq. (15) with $s(t)$ from Eq. (21).

Under the influence of the linear electromagnetic field, the model can present a positive expansion and negative deceleration simultaneously, then producing inflation, however, the period of inflation is shorter than the corresponding to the BI field. Moreover, the interpretation of the spacetime is not clearly cosmological for the periods when $s < 0$ ($t > 0.5$ in Fig. 4). The conclusion is that, for the same geometry, the linear field does not favour inflation as much as BI does. The effect is shown in Fig. 4.

VII. EQUATION OF STATE AND ENERGY CONDITIONS

We can ask about the behavior of the nonlinear radiation or the relationship between the energy density and the pressure. Such relation can not be the equation of state of a perfect fluid because the algebraic structure of




FIG. 6: It is shown the behavior of the matter with the pressure p as function of the energy density ρ . Both pressure and energy density are scaled as p/b^2 and ρ/b^2 . Also are shown, for comparison, the “vacuum-like” state of $p = -\rho$ as well as the equation of state of the radiation, $p = \rho/3$. This behavior occurs during the period that the spacetime is anisotropic.

the energy momentum tensor $T_{\mu\nu}$ in our problem is not compatible with that of a perfect fluid. To correspond to a perfect fluid $T_{\mu\nu}$ should have three repeated eigenvalues. Rather, in the studied case, the energy momentum tensor possesses two double repeated eigenvalues that for an anisotropic space they can be interpreted as a different behavior of matter depending on the direction. The two double repeated eigenvalues of $T_{\mu\nu}$ are

$$T_{12} = 2b^2(e^\nu - 1), \quad T_{34} = 2b^2(e^{-\nu} - 1). \quad (23)$$

When the energy-momentum tensor has only one time-like eigenvector (E_4) one may express the components T^{ab} of the energy-momentum tensor at a point p^* with respect to an orthonormal basis in the canonical form $T^{ab} = \text{diag}[p_1, p_2, p_3, \rho]$, where ρ represents the energy density measured by an observer whose world line at p^* has as unit tangent vector E_4 ; the eigenvalues p_α represent the pressures in the three spatial directions. The corresponding eigenvalues are $\rho = p_1 = T_{34}$ and $p_2 = p_3 = T_{12}$, they are plotted in Fig. 5. One of the eigenvalues corresponds to the energy density and the pressure in one direction, while the other one corresponds to the pressure in the other two directions. The observer at p^* measures in one direction a stiff matter $\rho = p_1 > 0$ while in the other two directions the pressures are negative $p_2 = p_3 = p < 0$. Fig 5 also shows that as the time T grows, the eigenvalues tend to the same value but with negative signs, i. e. $\rho + 3p = 0$, it implies that the trace of the energy-momentum tensor becomes zero, and that the matter has the equation of state $p = -\frac{\rho}{3}$. Note that it coincides with the equation of state of the gas of Nambu-Goto strings in three spatial dimensions.

The energy density is given by

$$\rho = \frac{2b^2}{8\pi} \left\{ \sqrt{\frac{b^2 + D^2}{b^2 - \check{H}^2}} - 1 \right\}, \quad (24)$$

Eq. (24) shows that the growth of ρ with the fields $\check{H}, D$ is slower than that in the Maxwell theory, where $\rho \sim \check{H}^2 + D^2$. While the negative pressures are:

$$p_2 = p_3 = p = \frac{2b^2}{8\pi} \left\{ \sqrt{\frac{b^2 - \check{H}^2}{b^2 + D^2}} - 1 \right\}, \quad (25)$$

or in terms of ρ , for these two directions and scaling ρ/b^2 and p/b^2 , we have

$$p = -\frac{\rho}{\rho + 1}, \quad (26)$$

The behavior of $p(\rho)$ is illustrated in Fig. 6. In these two directions the matter behaves as a “vacuum-like” state. The observer at p^* sees that there is a co-existence of an expanding-contracting asymmetry: for $p = -\rho/(\rho + 1)$ produces expansion in two directions (negative pressures) whereas contraction prevails in the third direction, $p = \rho$, producing together a global expansion shown in Fig.1.

In the two directions of negative pressure, the compressibility is negative, $dp/d\rho < 0$; it is well known that negative compressibility leads to a strongly inhomogeneous mixture of different phases. From Eqs. (24) and (25), one has also that

$$\rho = -2b^2 p / (8\pi p + 2b^2), \quad (27)$$

This equation indicates that for values of the pressure such that $p = -\frac{2b^2}{8\pi}$ there occurs a critical point in the energy density, ρ diverges for this value of the pressure. It corresponds to the values of the fields that superate the maximum field strenght, $\check{H} \geq b$. The instability in the energy density indicates a possible phase transition at $\rho > \rho_{crit}$. It is known that Born-Infeld theory for values of the invariants much greater than the limit field, $P \gg b^2$, i.e. in the “super-quantum limit”, is equivalent to the theory of magnetic strings. Hence in spacetime regions of strong field the self-trapping of the field into condensate of closed Nambu strings should be expected [19].

In [6] Altshuler investigated nonlinear electrodynamics as an alternative way to inflation and the possibility of anti-inflation. Altshuler addressed the case of the Lagrangian depending in nonlinear way only on one of the invariants of the electromagnetic field; however some of the conclusions drawn there applies to our case for the directions of observed negative pressures. One of the conclusions of his study is that for sufficiently high density the radiation, governed by nonlinear dynamics, has

a tendency to condensation when the effective pressure is negative. This condensate state is the responsible for inflation. However, strictly speaking, to obtain an effective equation of state we should make a statistical approach, like a kind of average in the three directions.

A. Energy Conditions

In this subsection the energy conditions for the model are analyzed.

The dominant energy condition (DEC) states that *for every timelike vector, W_a , $T^{ab}W_aW_b \geq 0$ and $T^{ab}W_a$ is a nonspacelike vector* [14]. This condition means that to any observer the local energy density appears nonnegative and the local energy flow vector is non-spacelike. In the case of the Born-Infeld energy momentum tensor DEC is a requirement of a physically reasonably nonlinear structural function and its fulfillment is guaranteed by condition (iii) in Sec. II. For the timelike unit vector U^α DEC amounts to

$$T_{ab}U^aU^b = 2T_{34}U^3U^4 = 2b^2(e^{-\nu} - 1) \geq 0, \quad (28)$$

clearly DEC is fulfilled if $e^\nu = \sqrt{1 - \varphi^4} < 1$, which in fact is the case we addressed in this work ($\varphi^4 < 1$).

The strong energy condition (SEC), as stated in [20], says: *A spacetime satisfies the strong energy condition if $R_{ab}u^au^b \geq 0$, for all causal vectors u^a* . The same condition is called by Hawking as the *timelike convergence condition* [14].

In the studied case the traceless Ricci tensor $R_{\mu\nu}$ has two double repeated eigenvectors such that in the Segré classification $R_{\mu\nu}$ corresponds to A1[(11)(1,1)]. Actually the two nonvanishing components of the traceless Ricci tensor are

$$R_{12} = 2b^2(1 - e^\nu), \quad R_{34} = 2b^2(1 - e^{-\nu}). \quad (29)$$

For the BI field SEC amounts to $R_{ab}U^aU^b = R_{34} = 2b^2(1 - e^{-\nu}) < 0$, which clearly violates SEC and the reason is the existence of the negative pressures ($p_2 = p_3 < 0$). This fact must not disturb us, since inflationary models driven by scalar inflaton fields violate SEC during the inflationary epoch. The minimally coupled scalar field violates SEC and indeed curvature-coupled scalar field theory also violates SEC. This point has been discussed recently in [20] and [21].

VIII. FINAL REMARKS

In this paper we showed that the Born-Infeld nonlinear electromagnetic field induces inflation in anisotropic cosmological models that are conformal to several Bianchi spaces. Moreover, the e-folding number to obtain enough inflation to solve the horizon and flatness problems can

be provided by nonlinear electrodynamics with a fine tuning of the parameter b . Note that changing the value of b imply physically different nonlinear electrodynamic theories coupled to gravity, since the coupling constant is different. The asymptotic behavior of the kinematical parameters shows that the inflation occurs in a limited period of time and that the model isotropizes, since as the synchronous time goes to infinity the expansion, deceleration and shear tend to vanish.

In contrast, the linear electromagnetic field inflates the model in a very restricted period of time and the range of the coordinate t valid for a cosmological interpretation is also reduced.

The analysis of the nonlinear electromagnetic radiation in the anisotropic space exhibits expansion in two directions, associated to negative pressures that could originate condensated states which allow inflation. In the third direction the equation of state corresponds to stiff matter, this fact could indicate compression in such direction.

We also analyzed the energy conditions and it turned out that the strong energy condition SEC is violated by the BI field. The violation of this condition could have as a consequence the absence of an initial singularity, since the fulfilling of SEC is related with the convergence of neighbouring geodesics. However, in [22] it was obtained the result that inflationary spacetimes are not past complete, i.e. the corresponding geodesics are incomplete and then a singularity exists. Therefore we wonder if this spacetime is singular. It is work in progress if in the case we addressed the nonlinear electromagnetic field does prevent the existence of an initial singularity.

Being the nonlinear electromagnetic field consistent

with an inflationary scenario, it might be that the BI field or another field with a Lagrangian of this type was present at early epochs of the universe, probably not alone, but accompanied with some scalar fields and that together, in a kind of “assisted inflation”, cooperate to produce the appropriate initial conditions for a FRW-type universe. However to address the standard picture of inflation the question arises on How can we consider quantum fluctuations of the BI field during inflation?, since in fact BI theory corresponds to an effective modification induced by quantum fluctuations. Also we must investigate if within the BI theory is it possible to have reheating in the universe; this lead us to the question of creation of particles and quantization of the BI field; this study was initiated by Born and Infeld themselves in 1935 and they showed that the commutation rules for the field components were very similar to the ones of quantum mechanics. Alternatively, one could adopt a scheme like the proposed in higher-derivative gravity theories and to look for a conformal equivalence between BI field coupled to gravity on one side and the Einstein theory with a scalar field on the other side, and then proceed as in the standard inflationary treatments. All these are speculations that have to be worked out.

Acknowledgments

The authors acknowledge one of the referees for a careful reading of our manuscript and for valuable suggestions. R. G-S. gratefully acknowledges fruitful discussions with J. M. M. Senovilla. R. G-S. was partially supported by FC-UAEMex. This work was partially supported by CONACyT (México) under project 40888A-1.

-
- [1] M. Born and L. Infeld, Proc. Royal Soc. London **A144**, (1934) 425
 - [2] W. Heisenberg and H. Euler, Z. Phys. **98** (1936) 714.
 - [3] P. Vargas Moniz, Phys. Rev. **D66**, (2002) 103501; Class. Quant. Grav. **19** (2002) L127-L134.
 - [4] A. Guth, Phys. Rev. **D23** (1981) 347.
 - [5] P. de Bernardis *et al*, Nature **404** (2000) 955.
 - [6] B. L. Altshuler, Class. and Quantum Grav. **7** (1990) 189.
 - [7] L. Jensen and J. Stein-Schabes, Phys. Rev. **D35** (1987) 1146; M. Demianski, R. de Ritis, G. Platania, P. Scudellaro and C. Stornaiolo, Phys. Rev. **D35** (1987) 1181.
 - [8] J. L. Cervantes-Cota, Class. Quant. Grav. **16** (1999) 3903.
 - [9] J. M. Aguirregabiria, A. Chamorro, L. P. Chimento, and N. Zuccalá, Phys. Rev. **D62** (2000) 084029.
 - [10] J. M. Aguirregabiria, P. Labraga and R. Lazkoz, Gen. Rel. Grav. **34** (2002) 341.
 - [11] R. García-Salcedo and N. Bretón, Int. J. Mod. Phys., **A15**, (2000), 4341.
 - [12] V. A. De Lorenci, R. Klippert, M. Novello, J. M. Salim, Phys. Rev. **D65**, (2002) 063501.
 - [13] V. V. Dyadichev, D. V. Galtsov, A. G. Zorin, M. Yu. Zotov, Phys. Rev. **D65**, (2002) 084007.
 - [14] S. W. Hawking and G. F. R. Ellis, *The large scale structure of spacetime* (Cambridge Univ. Press, 1973.)
 - [15] J. F. Plebanski, *Lectures of Nonlinear Electrodynamics* monograph of the Niels Bohr Institute Nordita, Copenhagen (1968).
 - [16] A. García, H. Salazar and J. F. Plebański, Nuovo Cim. **84** (1984) 65.
 - [17] D. Kramer, H. Stephani, M. A. H. MacCallum and E. Herlt, *Exact Solutions of Einstein's Field Equations*, (Cambridge University Press, Cambridge, 1980.)
 - [18] C. B. Collins and S. W. Hawking, Astrophys. J. **180** (1973), 317.
 - [19] H. C. Tze, Nuovo Cimento **A22** (1974) 507
 - [20] J. M. M. Senovilla, Gen. Rel. Grav., **30** (1998) 701-848
 - [21] M. Visser and C. Barceló, *Energy conditions and their cosmological implications*, gr-qc/0001099.
 - [22] A. Borde, A. H. Guth and A. Vilenkin, Phys. Rev. Lett. **90**, (2003) 151301.